

| DEPTH | | C O R E | LITHOLOGY | EPOCH | STAGE | AGE (ZONE) | BIOSTRATIGRAPHIC NOTES |
|-------|-----|------------------|---------------------------------------|----------------------|---------|------------|---|
| m | ft | | | | | | |
| | | D1 | | Pleistocene | | | |
| | | C1 | | | | | |
| | | D2 | | Pliocene-Pleistocene | | | Ages based on Foraminifera and nannofossils Radiolaria absent. |
| | | C2R | | | | | |
| | | D3 | | | | | |
| | | D4 | | Pliocene | Upper | | |
| | 100 | C3 | | | | | |
| | | D4 | | Pliocene | Mid-Lwr | | |
| | | D4 | Clay Qtz Paly Kao Mica | | | | |
| | | C5 | Clay Qtz. Kao. Cris. Paly. sep. | | | | |
| | | C6 | | | | | |
| | 200 | C7 | Clay Qtz. Kao. Phil sep. Paly | | | | |
| | | C8 | Clay Qtz. Cris. Kao Paly | | | | Barren |
| | | C9 | Clay Paly Sep. Qtz Cris. | | | | |
| | 300 | C10 | Clay Paly Kao qtz | | | | |
| | | C11 | Clay Paly Kao | | | | |
| | 100 | C12 | Clay Paly Sep. Qtz. | | | | |
| | 400 | B1 | Clay Paly Qtz. | | | | |
| | | | | | | | |
| | 500 | B2 | Clay Paly. Cris. | Eocene | | | Radiolarian fauna sparse and poorly preserved. |
| | | | | | | | |
| | 600 | | | | | | |
| | 200 | | | | | | |
| | 700 | B3 | Clay Paly. Sep. Cris. Dol. | | | | |

STRATIGRAPHIC SUMMARY CHART - LEG 2 SITE 12
 Lat. 19°41.73'N Long. 26°00.03'W Water depth 4557 m
 Depth of penetration 215 m (Hole 12B)
 114.9 m (Hole 12C)
 36.6 m (Hole 12D)