

18. DISTRIBUTION OF FIRST SERIES TRANSITION METALS IN ROCKS RECOVERED DURING DSDP LEG 22 IN THE NORTHEASTERN INDIAN OCEAN¹

Henri Bougault, Centre Océanologique de Bretagne, Brest, France

INTRODUCTION

Ti, V, Cr, Fe, Co, Ni, Cu, and Zn have been analyzed from Sites 211, 213, 214, 215, and 216 of DSDP Leg 22 of the *Glomar Challenger* in the northeastern Indian Ocean. According to these trace elements, three different types of source material are broadly defined: the basin floor basalts with a composition similar to tholeiites from mid-oceanic ridges, the basalts and the differentiated rocks from the Ninetyeast Ridge, and the diabase and amphibolite from Site 211.

In this paper the notion of stabilization energy in a ligands field is summarized, analytical techniques used are described, and certain conclusions based on the results are presented.

STABILIZATION OF TRANSITION ELEMENTS IN A LIGANDS FIELD

The elements of transition of the first series differ from each other by the filling of their 3d energy levels; 3d orbitals of corresponding ions are partially filled. All the 3d orbitals have the same energy, but ions are surrounded by other anions or dipoles and it can be demonstrated that the five 3d orbitals must be divided into two types of orbitals: two orbitals "eg" and three orbitals "t_{2g}" possessing different energies (Orgel, 1964). Thus, the energy of an ion of a transition metal is a function of the number of electrons present in each orbital; this energy, which may be a few kilocalories per mole, has an important effect on the geochemical behavior of the transition metals (Burns et al., 1964; Curtis, 1964; Wagner and Mitchel, 1951). The theoretical variations of this stabilization energy related to an octahedric configuration in function of the atomic number is shown in Figure 1.

Allegre et al. (1968) proposed the use of the diagram log (concentration of the element in rock/concentration of the element in chondrite) for the transition elements versus the atomic number of the element, which was commonly used for rare earth studies. An example of the diagram is presented in Figure 2. The effect of the stabilization energy for the different metals is clearly shown (the W-shaped plot); the W shape of the diagram is increasingly sharp as the magmatic differentiation is more pronounced.

ANALYTICAL METHODS

Ti and Fe were analyzed by a classic X-ray fluorescence method using a glass disk made of 6 g of Li₂B₄O₇, 1 g of La₂O₃, and 1 g of calcinated powdered sample; G 2, BCR

1, PCC 1, DTS 1, PCC 1, and AGR 1 have been used as standards for calibration curves. Cr, Ni, and Co determinations have been obtained both by X-ray fluorescence using pellets and by atomic absorption. The solutions used for atomic absorption analyses were obtained with HF-H₂SO₄ dissolution of the powdered sample, evaporation to dryness, and dissolution with 50 cm³ of NCl N. Calibration curves were obtained with solutions containing the same quantities of major elements as found in the rocks. A C₂H₂-N₂O flame was used for V, Cr, Ni, and Co determinations; C₂H₂ air flame was used for Cu and Zn.

RESULTS

The results of the trace element analyses are shown in Table 1. All concentrations have been expressed in ppm metal; at least two samples per site were analyzed. The two samples of Site 211 represent two different types of rocks; all other results were obtained from two different samples of the same rock type.

The diagram log (concentration of the element in rock/concentration of the element in chondrites) for Site 211 is shown in Figure 3, for Sites 213 and 215 in Figure 4, and for Sites 214 and 216 in Figure 5. All the analyzed rocks plot on a W-shaped diagram. The values of element concentrations in chondrite chosen for calculating the ratio are as follows: Ti: 660; V: 50; Cr: 2700; Mn: 2500; Fe: 250,700; Co: 800; Ni: 13400; Cu: 100; Zn: 50.

Two pairs of elements are considered in Figures 6 and 7. The V-Cu pair have low stabilization energy and Cr-Ni have a high stabilization energy. The combined effects of the partition coefficients of these elements directly related to their stabilization energy, together with the magmatic differentiation, give a positive correlation between V and Cu on one hand and between Cr and Ni on the other. The correlation found for these two pairs of elements suggests that the notion of stabilization energy due to ligands field can be applied to magmatic processes.

The behavior and the values of these transition elements leads us to group the different sites as follows: Sites 214 and 216 from the Ninetyeast Ridge, Sites 213 and 215 from the Indian Ocean basin floor, and Site 211 situated 300 km west of the Java Trench.

Site 211: The sample analyzed from Site 211, Core 12, Section 1, 143-145 cm is a diabase while the sample from Core 15, Section 2, 0-5 cm is an amphibolite. Except for Ti, the abundances of the considered trace elements are higher in the diabase sill than in the amphibolite; this is very significant for Ni (127 and 75 ppm) and Cr (275 and 70 ppm). From the partition coefficient data it was suggested (Schwarcz, 1967) that the stabilization energy Δ is higher for biotite than for hornblende. Since the content of biotite in the diabase sill is higher than that of amphibolite (Chapter 17), it is likely that the abundances of Cr and Ni

¹Contribution No. 164 of the Département Scientifique, Centre Océanologique de Bretagne.




Figure 1. Stabilization energy in an octahedral field.

in these rocks are related to the content of minerals associated with a differentiation process.

Sites 213 and 215: From Site 213 one sample was analyzed from Core 18, Section 1 at 144-150 cm and one from Section 2, 101-103 cm. From Site 215 one sample was analyzed from Core 18, Section 2, 106-110 cm and one from Core 19, Section 2, 145-150 cm.

Using the McDonald and Katsura (1964) silica versus total alkalis variation diagram, it is observed that these samples from Sites 213 and 215 have very near the Hawaiian tholeiite and alkali basalt division line with an equation of: $Y = 0.37X - 14.4$ ($X = \text{wt\% SiO}_2$ and $Y = \text{wt\% Na}_2\text{O} + \text{K}_2\text{O}$). Little is known about the distribution of the transitional metals in alkali basalts and tholeiites. From the present data, it is observed that the transition metal distribution in the rocks of these two sites is very close to the mean distribution of mid-oceanic ridge basalts.

Sites 214 and 216: Site 214 contained two types of igneous rocks: a differentiated flow (oceanic andesite, Chapter 17) with an SiO_2 content higher than 54% and a basaltic flow. Basalts from both Sites 214 and 216 have a similar transition metal distribution. Table 1 and Figure 5 show high values of the ratio (element in rock/element in chondrite) for Ti-V and for the Cu-Zn pairs. The Ni-Cr

contents are lower than in mid-oceanic ridge basalts. These results suggest that differentiation has given rise to the distribution of the trace elements. There is a high concentration of elements with low stabilization energy and a low concentration of elements with high stabilization energy. It is difficult to classify the basalts from Sites 214 and 216 as tholeiitic or alkali basalts. Nevertheless, the rocks from these two sites on the Ninetyeast Ridge are different from tholeiites of the mid-oceanic ridges as far as the transition metals are concerned:

	Ti	V	Cr	Fe	Co	Ni	Cu	Zn
Ninetyeast Ridge	13,000	420	60	95,000	60	45	120	100
Tholeiite	7,000	250	230	81,550	60	100	60	60

Two samples of differentiated rocks from Site 214 were also analyzed. They are located in Core 48, Section 1, 94-100 cm, from a transition zone, and in Core 49, Section 1, 137-144 cm. From a transition metal distribution point of view, they are the most differentiated rocks (Figure 5). The concentrations of Ni, Cr, and V are very low (Table 1).




Figure 2. Evolution of the transitional elements in the Pindo amphibolitic suite : chondrite-normed diagram.

TABLE 1
Transitional Trace Element Analyses of Rocks Recovered from DSDP Leg 22 in the Indian Ocean

Site	Sample	Ti	V	Cr	Fe	Co	Ni	Cu	Zn	Rock Type
211	211-15-2,0-5	15,100	115	70	63,600	36	75	21	70	Amphibolite Diabase
	211-12-1,143-145	13,280	140	275	73,800	69	127	31	80	
214	214-53-1,26-30	12,700	390	80	103,000	59	55	31	104	Basalt Ninetyeast Ridge
	214-53-1,30-35	12,100	385	75	110,000	60	49	111	104	
	214-53-1,97-100	12,100	400	77	104,700	60	50	110	100	
	214-53-1,28-30	13,500	420	57	92,500	62	39	165	106	
	214-54-2,117-127	12,800	370	62	93,000	60	45	153	91	
216	216-37-2,80	16,570	420	50	90,700	60	33	147	100	Basalt Ninetyeast Ridge
	216-38-2,145-150	16,300	420	50	98,900	60	27	122	100	
214	214-48-1,94-100	8,340	38	4	70,400	33	10	5	115	Differentiated rocks Ninetyeast Ridge
	214-49-1,137-144	8,760	38	0	68,880	44	5	5	115	
213	214-18-1,144-150	6,060	235	280	74,000	68	81	80	75	Basalt
	213-18-2,101-103	6,180	235	290	73,800	66	85	80	85	
215	215-18-2,106-110		200	200		50	92	47	55	Basalt
	215-19-2,145-150	10,000	220	200	62,300	50	91	75	55	

Figure 2, which shows the ratio of the distribution of trace elements in various types of rocks, indicates that the concentrations of V and Cu increases from a peridotite to a basalt and then decreases slightly from a basalt to an albitophyre. Similar observations about the behavior of V and Cu were mentioned by Siedner (1965) for the fractionation of an alkali igneous suite. Although no ultrabasic samples are available from the Indian Ocean sites, the evolution of V and Cu can still be observed. The basalts from the Ninetyeast Ridge sites (214, 216) with relatively high V and Cu content (400 and 120 ppm, respectively) are more differentiated than those from the basin floor sites (213, 215) having a V content of 200 ppm and a Cu content of 80 ppm. The intermediate differentiated rocks from Site 214 have low concentrations of V and Cu (38 and 5 ppm, respectively).

CONCLUSIONS

The distribution of the transitional trace elements suggests the following conclusions:

1) The observed difference between the diabase sill and amphibolite from Site 211 is due to the mineralogical differences between these two rocks.

2) The basalts from Sites 213 and 215 situated in the central Indian Basin and in the Wharton Basin on both sides of the Ninetyeast Ridge are similar to mid-oceanic ridge basalts.

3) The basalts from Sites 214 and 216 of the Ninetyeast Ridge are different from tholeiites of mid-oceanic ridges. The differentiated rocks from Site 214 are the most

differentiated type of sample studied here as far as the transitional trace elements are concerned.

REFERENCES

- Allegre, C. J., Javoy, M., and Michard, G., 1968. Etude de l'abondance des éléments de transition dans l'écorce terrestre, comparée à celle des terres rares: *In* *Originie et distribution des éléments*. Ahrens, L. H. (Ed.). Pergamon Press, p. 914-928.
- Burns, R. G., Clark, R. H., and Fyfe, W. S., 1964. Crystal field theory and some geochemical applications: *In* *Chemistry of the earth's crust*, Vernadsky Centennial Symp., v. 2, p. 88-106.
- Curtis, C. D., 1964. Applications of the crystal-field theory to the inclusion of trace transition elements in minerals during magmatic differentiation: *Geochem. Cosmochim. Acta*, v. 28, p. 389-403.
- McDonald, G. A. and Katsura, T., 1964. Chemical composition of Hawaiian lavas: *J. Petrol.*, v. 5, p. 82.
- Orgel, L. E., 1964. *Chimie des éléments de transition*: Dunod Ed., Paris.
- Schwarz, H. P., 1967. The effect of crystal field stabilization of transition metals between metamorphic minerals: *Geochim. Cosmochim. Acta*, v. 31, p. 503-517.
- Siedner, G., 1965. Geochemical features of a strongly fractionated igneous suite: *Geochim. Cosmochim. Acta*, v. 29, p. 113-137.
- Wagner, L. R. and Mitchell, R. L., 1951. The distribution of trace elements during strong fractionation of basic magma: A further study of the Skaergaard intrusion, East Greenland: *Geochim. Cosmochim. Acta*, v. 1, p. 121-208.




Figure 3. Site 211: chondrite-normed diagram.




Figure 4. Sites 213 and 215: chondrite-normed diagram.




Figure 5. Sites 214 and 216: chondrite-normed diagram.




Figure 6. V-Cu correlation, diagram of two elements with low stabilization energy.




Figure 7. Cr-Ni correlation, diagram of two elements with high stabilization energy.