

INDEX

- Abyssal region, foram taxa in, 780
Accreted wedge, 572, 939, 1255, 1299
 definition of, 6
Accumulation rates, calculation of, 531
Active convergent margin, 809
Active oceanic ridges, 1259, 1263
Aleutian arc, Cenozoic volcanism on, 1030
Aleutian Trench, 1337
Aliphatic alcohols, 1377
 hydrocarbons, 1369
 ketones, 1375
Alkalinity, Japan Trench sediments, 1270
Aluminum/titanium ratio, 1219
Ammonia, Japan Trench sediments, 1270
Amorphous constituents in sediments, 941
"Amorphous" silica, 1214, 1229
Analcime, detrital, 1151
Andesite boulder, radiometric dating of, 1281
Andesite pebbles, 1090
Angular unconformity, 476, 490, 494, 560
Ankerite, 1123
Anoxic conditions, 553
Antarctic Circumpolar Current, 668
Anticline, breached, Japan Shelf, 1270
 $^{40}\text{Ar}/^{39}\text{Ar}$ ages of igneous rock, 1281
Arc volcanism, 24
Argillaceous sediments, 1233
Aromatic hydrocarbons, 1296, 1372
Aseismic zone, hypothetical convection cell in the,
 1296, 1372
Ash beds, alteration and diagenesis, 1030
Ash layers, distribution of, 1031
 graded, 545
 reworking of, 37
Ash SEE ALSO: Volcanic ash
Aso Volcano, 1287
Athabasca Tar Sands, 1344
Authigenic constituents, 1264
Authigenic dolomite, 1123
Back-arc basin, 569
Backbone Range Orogeny, 782
Basalt, East Pacific Rise, 1286
Basalt pebble, 1091
Bathymetry, benthic foraminifera, 780
Benioff Zone, 3
Bering Sea, Volcanic ash recovered from, 1064
Biogenic calcite, 1123
Biogenic methane, 1335
Biogenic opal, 1213
Biogenic silica, 542
 accumulation of, 542, 1229
 preservation and diagenesis of, 543
Biostratigraphic correlation of the Japan Trench
 Transect sections, 520 Figure 12
Biostratigraphic zonations, 6, 8–9 Figure 3
Biostratigraphy, Hole 438A, 511 Figure 4
 Hole 438B, 512 Figure 5
 Japan Trench Transect, 505
 Nannofossils, 875
Biotite, chemical composition of, 1061
Biotite hornfels, 1092
Bioturbation, 911
Biscuit structure, 576, 577
Bitumens, 1329
Bitumoids, 1331
Black shale, upper Cretaceous, 7, 553, 818, 1222
Bonin Trench, 467
Boron, 1225
Bouguer gravity anomalies, Japan Trench area, 454
 Figure 5
Breccia, defined, 576
Brunhes magnetic polarity epoch, 778
Brunhes/Matuyama Paleomagnetic Epoch boundary,
 197
Bulk density profiles, 582
Burial diagenesis, 1126
Burrows, 921
Butane, 1339
 $\text{C}_2\text{-}\text{C}_5$ alkanes, origin of, 1344
Calc-alkaline magmas, 502
Calcite, 1121
Calcite, biogenic, 1123
Calcium, 1223
Capistrano Formation, 666
Carbon, geochemistry of, 1319
 isotopic composition of, 1313
 stable isotopes of, 1285
Carbon-carbonate analysis, method of, 19
Carbon preference index (CPI), 1296, 1332
Carbonate compensation depth, 780
Carbonate constituents, origin of, 1120
Carbonate content of sediments, 941
Carbonate, diagenesis, 1123
Carbonate minerals, chemical composition of, 1118
 occurrence in cores, 1118
 X-ray diffraction studies, 1117
Carbonate nodules, 1094
 chemical composition of, 1095
Carbonate sediments, microtextures in, 1118, 1119
 scanning electron microscope studies of, 1119
 types of, 1120
Carboxylic acids, 1367
Cariaco Trench, 1337, 1395
Cartenoids, 1367, 1383, 1395
Channel filling, 556
Chert, defined, 1133
Cherts and procellanites, physical properties of, 1137
Chlorinity decrease downhole, Japan Shelf, 1270

- Chlorins, 1395, 1397
 altered, 1404
 Chlorite, 983
 Chlorophyll diagenesis, 1391, 1397
 Chondrites, 912
 Choshi Sequence, 687
 Chromium, 1225
 Circum-Pacific mineral megaprovince, 1006
 Clay minerals, 941
 authigenic origin of, 987
 composition of, 1012 Table 1
 electron diffraction analysis and electron microscopy, 979
 fluctuation of content, 535
 paleogeographic and diagenetic aspects, 979
 representative DTA-TG curves, 1014
 terrigenous origin of, 987
 Climatic conditions, cool and temperate, 897
 Climatic cooling, 198
 Climatic oscillations, 811, 821
 Clinoptilolite, 1145
 scanning electron microscope studies, 1151
 Cobalt, 1224
 Coiling direction history, *N. pachyderma*, 778
 Conductivity values, correlation of, 15
 Consolidation characteristics of sediments, Site 436, 1201
 Site 438, 1201
 Site 440, 1201
 Continental crust, Japan, 494
 Continental shelf, rapid subsidence of, 1270
 Convergent margins, 3, 473, 475, 569
 nonaccreting, 487
 Cool climatic intervals, North Pacific, 821
 Cooling event, upper Miocene, 773
 Copper, sediment content of, 1223, 1251
 Core disturbance, 585
 Cores, handling of, 17
 Cretaceous erosional surface, 193
 Cretaceous shales, 7, 553, 818, 1222
 maturity of, 553
 Cretaceous unconformity, 4
 Curie Temperature, 1079
 Cyclohexene, 1353
 Dacite boulder, radiometric dating of, 1281
 Dacite boulder conglomerate, 1083, 1126, 1285
 chemical composition of, 1084, 1086
 petrology and chemistry of, 1083
 potassium-argon age, 1084
 Dacite pebbles, 1090
 Deep sea terrace, 489, 494
 subsidence of, 482, 564
 Deposition mechanisms, deep ocean basin, 558
 Detrital calcite, 1123
 Detrital constituents, 1264
 Detrital sediments, Japan Trench, 1067
 Dewatering, structures formed by, 486
 Dewatering of sediments, 468
 Diagenesis, 550, 1143, 1144, 1151, 1384
 of clay minerals, 1015
 organic matter, 552, 1305
 sediments, role of clay minerals in, 993
 Diagenetic transformations, 1367
 Diasterenes, 1371
 Diatoms, basis for zonation of, 663
 biostratigraphy of, 633, 646
 correlation with onshore sequences, 687
 correlation with paleomagnetic stratigraphy, 665
 from Sites 436, 438, 440, correlation with Choshi section, 688 Figure 2
 correlation with Joban and Fukushima-Miyagi section, 689
 correlation with Sanno-ha-Kadonosawa section, 691
 correlation with Toyokoro sections, 692
 Hole 434, 636 Table 2
 Hole 436, 638
 Hole 438, 646, 647 Table 1
 Hole 438A, 649 Table 2, 650 Table 3, 651 Table 4, 652 Table 5, 653 Table 6, 646 Table 7, 655 Table 8
 Stratigraphic range of selected, 656 Figure 4
 Hole 438B, 655, 658 Table 9
 New variation, *Thalassiosira nidulus* var. *delicata*, 671, 684 Figures 1, 4
 preservation of at subduction zones, 1273
 Site 434, 362
 Site 435, 197, 635
 Site 436, 406
 Site 438, 39, 646
 Site 439, 656
 Site 440, 231, 659 Table 10, 660 Table 11, 661 Table 12
 Site 441, 328, 664 Table 13
 taxa, 634 Table 1
 zonation, 505, 635, 647 Figure 3
 defined, 641
 Dissolution of forams, relationship to climatic cool periods, 821
 Dolomite, 1121
 association with ankerite, 1270
 authigenic, 1123
 occurrence in carbonate sediments, 1118
 Downhole logs, Leg 57, 1169
 Sites 438/439, 1171
 Site 440, 1173
 Site 441, 1180
 Down-slope mass movement, 486, 1298
 Earthquakes, frequency-magnitude distribution, 468
 hypocenters of, 467
 East Pacific Rise, basalt from, 1286
 Elements, accumulation rates of, 1264
Emiliana huxleyi, lipid extract of, 1374
 Epoch boundaries, defined, 645
 Erratic pebbles, petrology of, 196
 Ethane, 1339
 Ethers, 1373
 “Excess” iron, 1220
 “Excess” silica, 1213, 1229
 origin of, 1234
 Explosive volcanism, history of on Honshu, 545
 pattern of, 1031
 rate of, 1034
 Japan Trench Transect, 1027
 Fatty acids, 1381
 Fault and block structures, 457

- Faults, 499, 500, 553
 defined, 576
 synsedimentary, 495
- Fecal pellets, abundance of, 922 Figure 1
 organic geochemistry, 923
 Site 436, 407, 921
 Type A, 922
 Type B, 922
 Type C, 922
 Type D, 922
- Fish teeth SEE: Ichthyoliths
- Flysch deposits, 1123
- Folding, 457
 synsedimentary, 1160
- Foraminifera, benthic, abyssal, 780
 bathymetry of, 780
 correlation with land section, 835
 correlation with other marine sections, 835
 Japan Trench area, 835
 Hole 438, 816 Table 4, 837, 838–839 Table 2, 840 Figure 2
 Hole 438A, 837, 842–845 Table 3, 846 Figure 3
 Hole 438B, 841, 842–845 Table 3
 Hole 439, 847
 Miocene, 841
 Site 435, 197, 786–787 Table 4
 Sites 438/439, 853 Table 5
 Site 439, 841
 Site 440, 847, 848–849 Table 4, 850 Figure 5
 Upper Cretaceous, 846
 upper Oligocene, 841
- Foraminifera, planktonic, biostratigraphy of, 776
 correlation with onshore sequences, 822
Globorotalia wilesi, new species, 785, 796 Figures 1–3
 Hole 438, 813
 Hole 438A, 46, 813, 814–815 Table 3
 Japan Trench, 775
 mid-latitude faunas in the North Pacific, 811
 modern lysocline, 780
N. pachyderma, coiling of, 821
 notes on species, 821
 Site 434, 364, 783, 784
 Site 435, 779, 783
 Site 436, 785, 407, 788 Table 5
 Site 438, 46, 812
 Site 439, 46, 818
 Site 440, 234, 818, 819 Table 5
- Fore-arc region, bathymetric map of, 523 Figure 1
 terminology of, 6 Figure 2
- Fractures, 553
 defined, 576
- Fractures and microfaults, healed, 577
- Fracturing, 4
- Free-base porphyrins, 1397, 1405
- Free lipids, 1367
- Fukushima-Miyagi sequence, 690,
- Gas chromatography (GC), 1292, 1327, 1369
- Gauss Magnetic Epoch, 232, 1160
- Geochemistry of trench sediments, 1259
- Geochronologic scale, 7
- Geologic setting, Japan Trench Transect, 522
- Geomagnetic data, 465
- Geosynclinal sequence, 473
- Geothermal gradient, 1030, 1151, 1298
- Geothermal observations, Japan Trench Transect, 480, 1207
- Gilbert Paleomagnetic Epoch, 505, 645
- Glass SEE: Volcanic glass
- Glaucite, 1019
- Global cooling, 667
- Global plate tectonic reconstruction, 483
- Graded bedding, 545
- Grain-size analysis, method of, 19
- Gravity measurements along the Japan Trench, 449
- Gravity transport, near-bottom, 1008
- Great Barrier Reef, ooids from, 1121
- “Green Tuff” tectonic belt, 473, 482, 1033, 1282
- Groundwater recharge, Japan shelf, 1270
- Hachinohe Knoll, 495, 522
- Hachinohe pumic, 1049
- Hadal region, benthic foraminifera taxa in, 780
- Hakobuchi Group, 1157
- Heat flow, effect on humic acid diagenesis, 1307
 thermal gradients, 480
- Heavy minerals, 1019
 Japan Trench sediments, 1020 Table 1
- Hemipelagic sediments, classification of, 11, 12 Figure 5
- Heptane, 1350
- Hidaka Basin, 34
- Hidaka Channel, 479
- Hidaka metamorphic belt, 474
- Hidaka Orogeny, 474
- Hidaka Trough, 474
- Hokkaido, Late Cenozoic sediments on, 1009
 metamorphic belts of, 464
- Hole 434, Radiolaria, 696
- Hole 434A, Radiolaria, 697
- Hole 436, correlation with land section, 772
 pollen analysis, 903
 pollen diagram, 904–905 Figure 1
- Hole 438, Foraminifera, 813, 816 Table 4, 837, 840 Figure 2
 Foraminifera, benthic, 838–839 Table 2
 Lithostratigraphy, 28
 Nannofossils, 877, 880 Table 3
 Paleobathymetry, 39 Figure 7
 Radiolaria, 736 Table 1
 sedimentation rates, 43 Figure 11
- Hole 438A, biostratigraphy, 511 Figure 4
 correlation with the land section, 772
 foraminifera, 46, 813, 814–815 Table 3, 842–845 Table 3, 846 Figure 3
 lithostratigraphy, 30
 nannofossils, 877, 882–883 Table 4
 paleobathymetry, 40 Figure 8, 511 Figure 4
 paleotemperature curve, 763 Figure 6
 pollen diagram, 900–901 Figure 2
 radiolaria, 738–743 Table 2
 sediment accumulation rate curve, 43 Figure 12, 657
- Hole 438B, biostratigraphy, 512 Figure 5
 diatoms, 658 Table 9
 foraminifera, 841, 842–845 Table 3

- nannofossils, 878, 884 Table 5
 paleobathymetry, 41 Figure 9, 512 Figure 5
 radiolaria, 697, 744 Table 3
 Holes 438A, 438B, and 439 lithostratigraphy, 31
 Hole 439, foraminifera, 847
 geothermal gradient in, 1015
 pollen diagram, 896–897 Figure 1
 radiolaria, 745 Table 4
 Hole 439 area, pre-Neogene proximity to a land source, 895
 Hole 440, radiolaria, 746 Table 5
 Hole 440A, radiolaria, 746 Table 6
 Hole 440B, radiolaria, 748–753 Table 7
 Honshu Island, 4
 volcanic front of, 1049
 Hornblende, 1069
 chemical composition of, 1061
 phenocrysts, 1076
 chemical composition of, 1077 Table 1
 Horst and graben structures, 476, 451
 Humic acids, 1307, 1367, 1383
 elemental composition of, 1308
 Humic organic matter, 895
 Huminite/vitrinite reflectance values, 1293
 Hydrocarbons, 1293, 1336–1353, 1367
 aliphatic, 1369, 1375, 1377
 aromatic, 1296, 1372
 Hydrogen sulfied odor, 1339, 1342
 Ice Age conditions, 772
 Ice-rafted material, 37
 Ice-rafting, 667, 1095
 Ichthyoliths, description of, 887
 preservation of, 887
 Site 436, 887, 888 Table 1
 Igneous rocks, magnetic susceptibility, 482
 Imbricate faulting, 4
 Inner trench sediments, chemical characteristics of, 1255
 Insoluble organic matter, 552
 Interstitial water, Japan Trench, 1269
In situ temperatures, Japan Shelf, 1274
In situ water samples, Japan Trench, 1269
 “Inundation” type of trench, 469
 Iron, 1220
 Iron and manganese, precipitation of, 1243
 Iron, “excess”, 1220
 Ishikari-Hidaka Basin, 474, 522
 source of sediments, 554
 seaward transgression of, 483
 Ishikari tectonic belt, 473
 Island arc, 1021
 Island-arc andesitic belt, 1005
 volcaniclastic material from, 1008, 1009
 Island-arc evolution, 1034
 Island-arc system, 1285
 Island arcs along active margins, 1071, 1078
 Isobutane, 1343
 Isopentane, 1343
 Jamaican reefs, scalanohedral calcite cement from, 1121
 Japan and Kuril Trenches, juncture of, 557
 Japan arc-trench system
 Japan Convergent Margin, 569
 deformation along, 585
 geologic history and structure, 570
 tectonic history of seaward edge, 502
 tectonic processes and evolution of, 486
 Japan fore-arc region, 569, 570
 Japan slope sediments, comparison of sediments to other convergent margins, 583
 Japan subduction zone, 1083
 Japan Trench, 993
 crustal structure of, 463
 foraminifera, 775
 formation of, 457
 geological cross section of, 56 Figure 19
 heavy mineral analysis of sediments, 1020 Table 1
 proposed evolution of, 57 Figure 20
 sedimentation in, 478
 subduction-accretion complex, 484
 tectonic stress associated with plate convergence in, 584
 west slope sediments, origin of, 1246
 Japan Trench lower slope, seismic records from, 496
 Japan Trench sediments, alkalinity of, 1270
 ammonia in, 1270
 clay minerals in, 1011
 coarse-silt fraction mineralogy of, 1005, 1006 Table 1
 Japan Trench Transect, 570, 1011
 biostratigraphic correlation of sites, 520 Figure 12
 biostratigraphic summary of sites along, 505
 heavy minerals in sediments from, 1019
 magnetic anomaly profiles from, 455 Figure 6
 main objectives of drilling in, 4
 multichannel seismic reflection records, 489, 490, 501
 pre-site survey of, 449
 sedimentation rates for, 538 Figure 13
 sediment composition, 939
 summary of, 473
 Japan volcanic arc, 1086
 Japanese Island Arc, 957, 1027
 Jaramillo event, 1160
 Jōban coal-field sequence, 688
 Kabato landmass, 473
 Kado area, 1072
 Kadonosawa Formation, 690
 Kamuikotan-Hidaku-Tokoro tectonic belt, 474
 Kamuikotan metamorphic belt, 474
 Kamuikotan tectonic zone, 1157
 Kerogen, 1293, 1327
 Kitakami-Abukuma Mountains, 451
 Kitakami-Abukuma tectonic belt, 473
 Kitakami Basin, 34
 Kitakami Massif, 4, 473
 Kitakami Mountains, granitic rocks, 1076
 Kuji Group, 1062, 1151
 Kunitan Formation, 1072
 Kurile Trench, recent sediments from, 1009
 Kuril volcanic arc, 1086
 Kuroshio current, 476, 505, 993
 Kwanto Region, 868
 Lead, 1225
 Legs 56 and 57 sites, correlation of, 514
 lithology of, 525–527 Table 1
 Lipid extracts, 1374
 Lipids, 1367
 concentration of in sediments, 1319

- Lipid source, bacterial, 1386
 marine, 1386
 terrestrial, 1386
 Lithification trends, 580
 Lithologic classification, 7
 Lithostatic versus tectonic dewatering, 581
 Long-baseline refraction measurements, 465
 Lower Japan Trench slope, structure of, 483
 Lysocline, 780
M-discontinuity, 465
 Macrofossils, 36
 Magmatic-arc rocks, emplacement of, 1084
 Magnesium, 1223
 Magnetic anomalies, offshore of the Tohoku Arc, 449
 Magnetic anomaly patterns, 465
 Magnetic anomaly profiles, 455 Figure 6
 Magnetic quiet zone, 449
 Malaga core section, 772
 Manganese content, marlstones, 1234
 sediments, 1220, 1221
 Manganese nodules, 1335
 Mariana arc and trench complex, 487
 Marine sediments, diagenesis of, 1243
 Mass spectrometry, 1292, 1369
 Matuyama reversed polarity epoch, 778, 1160
 Matuyama/Gauss paleomagnetic epoch, 197
 Maurice Ewing Bank, 668
 Messinian Salinity Crisis, 42
 Metal accumulation rates, 1254, 1256
 Metal enrichment, 1251
 Metalliferous sediments, 1259, 1263
 Metalloporphyrins, 1397, 1405
 Metamorphic belts, 464, 474
 Methane, biogenic, 1335
 generation of, 553, 1335
 origin of, 1344
 Metogahana Formation, 687
 Microfaults, 553
 Microfossil zones, correlation with geologic time scale, 509
 Microfracturing, 486
 Mid-America Trench, 487
 Mid-slope terrace, 572
 definition of, 6
 Miocene/Oligocene boundary, defined, 505
 Miocene/Pliocene boundary, 232, 496, 500, 645, 772
 defined, 505
 Minato Formation, 1072
 Mixed-layer minerals, 1015
 Monterey Formation, 666
 Monterey shale, 1133
 Mt. Shikotsu pumice fall, 1049
 Moho discontinuity, 4
 Multichannel seismic reflection record, 482, 489, 580
 Japan Trench Transect, 490, 501
n-alkanes, 1332
 Na-ami Formation, 687
 Nankai Trough, 554
 Nannofossil assemblages, paleobiogeography, 869
 biostratigraphy of Japan Trench, 875
 dissolution of, 877
 Hole 438, 877, 880 Table 3
 Hole 438A, 877, 882–883 Table 4
 Hole 438B, 878, 884 Table 5
 Japan Trench upper slope 867
 Pliocene paleobiogeography of, 869 Table 2
 Site 435, 867, 868 Table 1
 Site 438, 45, 877
 Site 439, 879, 884 Table 5
 Site 440, 233, 879, 885 Table 6, 886 Table 7
 Site 441, 329, 880
 zonation of, 505, 875
 Needle probe measurements, 15
 Nemuro tectonic belt, 474
 Neogene/Cretaceous boundary, 500
 Neogene volcanic ashes, 1027
 Neogene volcanism, 1033
 Neopentane, 1337, 1343
 origin of, 1345
 Nickel content, 1224
 Noda Group, 1072, 1151
 Non-aromatic hydrocarbons, 1293
 Non-clay minerals in sediments, 941
 Normal faults, 553
 Northeast Japan Arc, 1062
 North Pacific ocean plate, movement of, 927
 Ocean Bottom Seismograph (OBS) refraction data, 459, 463, 489
 Oceanic crust, 496
 Offscraping, 1248
 Oga Peninsula, 1079
 Olduvai Paleomagnetic Event, 505, 646, 778, 1160
 Opal, biogenic, 1213
 Opal-CT, 1135, 1145, 1274
 Opaline silica, X-ray diffraction studies, 1149
 Organic carbon, 1327, 1331
 accumulation rate of, 552
 Legs 56 and 57, 555 Figure 25, 1293
 Organic geochemistry, Legs 56 and 57 sediments, 1327, 1367, 1391
 Organic matter, bituminological characteristics of, 1333
 diagenesis of, 552, 1305
 maturation of, 553, 1330
 Organic petrography, 1291
 Osaka Group, 1057, 1060
 Ostracodes, Site 435, 907
 taxonomic notes, 907
 Oxygen-minimum layer, 1298
 Oyashio Current, 198, 476, 505, 771
 southward displacement of, 7
 Oyashio and Kuroshio currents, mixing zone of, 810
 Oyashio landmass, 490, 991, 1009, 1067, 1071, 1075, 1078, 1157, 1281, 1282, 1298
 erosion surface on, 560
 subaerial extent of, 499
 subsidence of, 4, 499, 500, 993
P-wave velocity measurement, 460 Figures 2 and 3, 461 Figures 4 and 5
 Pacific diatom zones, 634 Figure 1
 Pacific margin, island arc volcanism on, 482
 recent sediments on, 1009
 Pacific Plate, 449, 465, 467

- advance of against the island arc mass, 468
 convergence of, 569
 subduction of, 939, 1047
Paleobathymetry, 849
 Hole 438, 39 Figure 7
 Hole 438A, 40 Figure 8, 511 Figure 4
 Hole 438B, 41 Figure 9, 512 Figure 5
Paleoclimatic interpretation, foraminifera, 821
Paleoclimatology, late Miocene, 669
 Pliocene, 669
 Quaternary, 669
Paleogeothermal gradient of the Japan fore-arc region, 1015
Paleomagnetic stratigraphy, 7
 Site 440, 1159
Paleotemperatures, 757, 763 Figure 6
Paleotemperature curve, Southern California, 761, 763 Figure 6
Palynology SEE: Pollen
 Pancho Rico Formation, 667
 Peedee Formation, 1286
 Pelagic clays, sedimentation history of, 991
 Pellet-armored rod, 912
 Peru-Chile Trench
 sediments in, 451
 Perylene, 1372
 Petroleum, generation potential of sediments, 1299
 sources of, 1328, 1332
 Physical properties, Legs 56 and 57, 615–629
 measurement of, 572
 methods, 12
 modification of by tectonism, 1198
 Site 434, 1193
 Site 435, 1190
 Site 436, 1195
 Sites 438/439, 572, 1187
 Site 440, 1190
 Site 441, 1193
 Trench inner slope sites, 574
 Phytenes, 1370
 Pillow basalts, 1134
 Plagioclase, chemical composition of, 1058
 phenocrysts of, 1075
 Plate convergence, 570
 Plate movement, 1248
 Plate subduction, mechanism of beneath Japan, 468
 Pliocene, environmental instability of, 869
 sedimentation rates during, 581
 Pliocene/Pleistocene boundary, 231, 234, 821
 defined, 505
 hiatus at, 4
 Pliocene-Pleistocene sedimentation, 564
 Pliocene-Pleistocene tectonics, 564
 Pliocene-Quaternary boundary, 645
 Pliocene tectonic event, 554
 Poisson's ratio, 461, 584
 Pollen, 1296, 1327
 Pollen analysis, Hole 436, 903
 Holes 439 and 438A, 895
 Pollen diagram, Hole 436, 904–905 Figure 1
 Hole 438A, 900–901 Figure 2
 Hole 439, 896–897 Figure 1
 Porcellanite, defined, 1133
 Pore pressure, 468
 Pore water, upward migration of, 486
 Potassium, 1223
 Pristane/phytane ratio, 1296, 1322
 Propane, 1339
 Pumpellyite, 196
 Pumpellyite metaquartzite, 1092
 Pyrite, authigenic, 1005, 1221
 Pyroxene, chemical composition of, 1058
 electron microprobe analyses of, 1044 Table 1
 Quartz, relation to temperature and geologic age, 1274
 Quick Thermal Meter (QTM), description of, 14
 Radiolaria, biostratigraphy of, 695, 737
 Collospheara pyloma, new species, 761, 768 Figures 5–9
 correlation of sites using, 755
 correlation with Japanese land section, 704
 Eucyrtidium asanoi, new species, 709, 726 Figures 12a–14b
 evolutional lineages, 697
 Hole 434, 696
 Hole 434A, 697
 Hole 434B, 697
 Hole 438, 736 Table 1
 Hole 438A, 738–743 Table 2
 Hole 438B, 744 Table 3
 Hole 439, 745 Table 4
 Hole 440, 746 Table 5
 Hole 440A, 746 Table 6
 Hole 440B, 748–753 Table 7
 reworking of, 756
 Site 434, 362, 696, 698–699 Table 2
 Site 435, 197, 697, 700 Table 3
 Site 436, 406, 697, 702–703 Table 4, 704 Table 5, 705 Figure 1
 Site 438, 44
 Site 439, 45
 Site 440, 233
 Site 441, 328, 754–755 Table 8
 Taxonomy of, 704, 761
 zones, correlation of between sites, 762
 defined, 737
 Radiolarian datums, 758–760 Table 9
 warm water, 771
 Radiolarian events, 701, 706
 Radiolarian zones, 505
 correlation of, 761 Figure 4, 772 Figure 2
 defined, 696
 Radiometric dating, 1281
 Rebun Group, 473
 Reducing zone, 1251
 Reflective sequence, definition of, 6
 Rhodochrosite, authigenic, 1005
 Rhyodacitic components, 1030, 1049
 Rhyolite boulder, radiometric dating of, 1281
 Rhyolite conglomerate, 1285
 Rhyolitic components, 1030, 1049
 Rikicho Formation, 490
 Rind burrow, 911
 Rosario Formation, 846
 S-wave velocity measurement, 461

- Salinity decrease with downhole depth, Japan shelf, 1269
 Sandstone pebbles, 1092
 Sandstone, provenance of, 1157
 Sannohe Group, 690
 Sannohe-Kadonosawa Sequence, 690
 Sanriku District, Honshu, 463
 Sea of Japan, opening of, 482
 Sediment consolidation studies in subduction zones, 570
 Sediment ponds, 457
 Sediment transport, 1008
 Sedimentary prism, 1251
 Sedimentary-tectonic evolution of continental margins, reconstruction of, 566
 Sedimentation, Japan Trench, 478 modes of, 1264
 Sedimentation rates, 1163, 1254, 1255
 Hole 438, 43 Figure 11
 Hole 438A, 43 Figure 12, 657
 Japan Trench Transect, 538 Figure 13
 nature of sediments as related to, 531
 reasons for variation between sites, 756, 771
 Site 440, 1265
 Sedimentation rate curve, 1165 Figure 6
 Site 440, 1164 Figure 4, 1265
 Sediments, chemical composition of, 1234, 1235–1238 Table 1
 Sediments, dewatering of, 468
 Seismic reflection profiling, 449
 Seismic refraction data, 459, 463, 489
 Seismic refraction measurements, 483
 Shallow-water environment, 36
 Shikoku subduction zone, 1109
 Shiratorigawa Group, 690
 Silica, 1213
 Silica, “amorphous”, 1214, 1229
 Silica, biogenic, 542, 543, 1229
 Silica diagenesis, Sites 438 and 439, 1143
 Silica, “excess”, 1213, 1229, 1234
 Silica lepispheres, tridymite, 1274
 Silica phases, relationship to temperature and geologic age, 1274
 Silver, 1225
 Single-channel seismic reflection surveys, 489, 500
 Sisquoc Formation, 666
 Site 434, lower Japan Trench slope, 355–398
 background, 355
 biostratigraphy, 362, 512, 518 Figure 10, 359
 carbonate nodule chips, 1089
 clay mineralogy and bulk composition of sediments, 944, 945–946 Table 4, 947 Figure 6
 clay mineralogy and related minerals, 981, 982 Table 3
 correlation of lithology with seismic reflection data, 366
 diatoms, 362
 exotic pebbles, 1089
 foraminifera, 364, 783, 784
 geochemistry, 366
 hydrocarbons, 1336
 lithification and deformation structures, 579
 lithology, 358, 530
 objectives, 356
 operations, 356
 petrology of pebbles, 361
 physical properties, 364
 principal results, 355
 radiolaria, 362, 696, 698–699 Table 2
 volatile organic components, 1353
 Site 435, Japan Trench upper slope, 193–224
 background, 193
 benthic foraminifera, 786–787 Table 4
 biostratigraphy, 197, 510, 514 Figure 7
 carbonate nodule fragments, 1090
 clay mineralogy and bulk composition of sediments, 944, 948–949 Table 5
 clay mineralogy and related minerals, 984, 985 Table 6
 diatoms, 197
 exotic pebbles, 1090
 foraminifera, 197, 779, 783
 geochemistry of sediments, 1253
 hydrocarbons, 1338
 inorganic geochemistry, 199
 lithification and deformation structures, 577
 lithostratigraphy, 194, 530
 nannofossils, 867, 868 Table 1
 objectives, 194
 operations, 194
 Ostracodes, 907
 paleobathymetry, 514 Figure 7
 physical properties, 198
 principal results, 193
 radiolaria, 197, 697, 700 Table 3
 sediment age, 868
 volatile organic compounds from, 1351
 Site 435 area, uplift of, 565
 Site 436, Japan Trench outer rise, 399–503
 Background, 401
 Biostratigraphy, 405 Figure 6, 406, 513, 519 Figure 11
 chemical trend, 1247
 clay mineralogy and bulk composition of sediments, 946, 951–953 Table 6
 clay mineralogy and related minerals, 984, 988–989 Table 11
 correlation of lithology with seismic reflection data, 400 Figure 2, 409
 diatoms, 406
 exotic pebbles, 1090
 fecal pellets, 407, 921
 foraminifera, 407, 785, 788 Table 5
 geochemistry, 409, 1253
 lithification and deformation structures, 577
 lithology, 404, 531
 objectives, 401
 operations, 403
 petrology of pebbles, 405
 physical properties, 407
 principal results, 399
 radiolaria, 406, 697, 702–703 Table 4, 704 Table 5, 705 Figure 1

- sedimentation rate, 479, 1278
 volatile organic compounds from, 1353
 Site 438, Japan Deep Sea Terrace, 23–192
 ash, 1027
 biostratigraphy, 507, 510 Figure 3, 511 Figure 4
 clay mineralogy and bulk composition of sediments, 946, 955–956 Table 7, 959 Figure 9, 960 Table 8
 carbonate diagenesis, 1117
 diatoms, 39
 foraminifera, 46, 812
 geochemistry, 1259
 hydrocarbons, 1339
 intervals of intense CaCO_3 dissolution, 813
 nannofossils, 45, 877
 operations, 24
 paleobathymetry, 510 Figure 3, 511 Figure 4
 primary objectives, 812
 radiolaria, 44
 sediment accumulation rates, 663
 sedimentation rate curves, 1165 Figure 6
 volatile organic compounds from, 1349
 Sites 438 and 439, background and objectives, 24
 bathymetric distribution of benthic foraminifera, 853 Table 5
 biostratigraphy, 37, 39 Figure 7, 40 Figure 8, 41 Figure 9, 42 Figure 10
 depositional environments and sedimentary evolution, 35
 geochemistry, 47
 lithification and deformation structures, 576
 lithologic units, 29 Table 2
 lithology compared, 522
 location of, 52 Figure 16
 logging, 51
 physical properties, 49
 rates of deposition, 36
 seismic reflection profiles, 51
 subsidence of, 560, 564
 summary and conclusions, 54
 Site 439, Japan Deep Sea Terrace, 23–192
 ash, 1028
 biostratigraphy, 507, 513 Figure 6
 carbonate diagenesis, 1117
 clay mineralogy and bulk composition of sediments, 949, 960 Figure 10, 961–963 Table 9
 dacite conglomerate, 481
 foraminifera, 46, 818, 841
 geochemistry, 1259
 hydrocarbons, 1341
 nannofossils, 879, 884 Table 5
 operations, 26
 paleobathymetry, 42, 513 Figure 6
 radiolaria, 45
 volatile organic compounds from, 1351
 Site 440, Japan Trench Midslope Terrace, 225–318
 ash, 1028
 background and objectives, 226
 biostratigraphy, 231, 232 Figure 5, 511, 515 Figure 8
 clay mineralogy and bulk composition of sediments, 949, 965 Figure 11, 966 Table 10
 conclusions, 241
 diatoms, 231, 659 Table 10, 660 Table 11, 661 Table 12
 foraminifera, 234, 818, 819 Table 5, 847, 848–849 Table 4, 850 Figure 5
 geochemistry, 234, 1262
 hydrocarbons, 1342
 lithification and deformation structures, 578
 lithostratigraphy, 226, 229 Figure 3, 530
 logging, 238
 nannofossils, 233, 879, 885 Table 6, 886 Table 7
 operations, 226
 paleobathymetry, 232 Figure 5, 515 Figure 8
 physical properties, 236
 principal results, 225
 radiolaria, 233
 sedimentation history and depositional environments, 231
 sedimentation rate, 663 Figure 7, 1164 Figure 4, 1265, 1342
 seismic reflection record, 237, 238 Figure 10
 stratigraphic ranges of diatoms and silicoflagellates, 662, Figure 6
 structural geology, 230
 summary, 240
 volatile organic compounds from, 1351
 Site 441 Japan Trench Lower Slope, 319–354
 background and objectives, 320
 biostratigraphy, 327, 329 Figure 10, 512, 516 Figure 9
 clay mineralogy and bulk composition of sediments, 950
 conclusions, 335
 diatoms, 328, 664 Table 13
 geochemistry, 330, 331 Table 2, 1263
 hydrocarbons, 1343
 lithification and deformation structures, 578
 lithostratigraphy, 321, 324 Table 1, 530
 logging, 333
 methane/ethane ratio, 1343
 nannofossils, 329, 880
 operations, 320
 physical properties, 331
 principal results, 319
 radiolaria, 328, 754–755 Table 8
 seismic reflection record, 331 Figure 12
 summary, 333
 Slickensides, 1107
 origin of, 1109
 Slump deposits, 584
 Slump scars, 476
 Slump structures, 476
 Slumping, 1095, 1328
 Slumps, rotational, 4
 Solid burrow, 912
 Sonobuoy measurements, Japan Trench, 465 Table 2
 Sonobuoy refraction data, 463, 464, 489
 Sonobuoy wide-angle reflection profile, 467
 Sorachi Group, 474, 1157
 Sterenes, 1295, 1370
 Stratigraphic hiatus, late Miocene, 42

- Upper Cretaceous/Neogene, 451
 upper Miocene, 232, 666
 Stratigraphy, Japan Trench, 476
 Subducted oceanic lithosphere, 1248
 Subduction, 569
 Subduction accretion, 569
 Subduction–erosion model, 486, 487 Figure 12
 Subduction mechanism acting on the Japan Trench, 469
 Subduction of ocean crust, 463
 Subduction zone sediments, geochemistry of, 1251
 Subduction zones, global variation of, 1033
 Northwest Pacific, 1034
 Submarine volcanoes, 1263
 Suenomatsuyama Formation, 690
 Sulfur, stable isotopes of, 1285
 Summary of results, Legs 56 and 57, 4
 Synsedimentary faulting, 495
 Synsedimentary folding, 1160
 Synthetic seismograms, 1180
 Taiki Formation, 691
 Takadate andesite, 1079
 Tarumae volcano, 1058
 Taup-Rotorua area ash, 1064
 Tectonic activity, 500
 Tectonic deformation, 578
 associated with convergence, 569
 of sediment, 553
 “Tectonic erosion” type of trench, 469
 Teichichnus, 912
 Tephra deposits, 1009
 Tephra layers, correlation of, 1045, 1058
 Tephras, petrographic properties of, 1043
 Tetrapyrrole pigment, 1392, 1395, 1401
 Thermal conductivity measurements, 480
 Thermal maturity, 1327
 Tin, 1225
 Tohoku Arc, frontal slope of, 449
 Tohoku Arc system, 457, 1062, 1064, 1079
 Tohoku Region, 1072
 Tokachi sequence, 691
 Tokoro belt, 474
 Toluene, 1350, 1353
 Tomesaki Formation, 690
 Torinosu limestone, 490
 Trace element distribution, sediments, 987
 Trace fossils, 911, 912
 Japan Trench, 911
 Transgressions, fore-arc basin, 483
 Transgressive Neogene sequence, 560
 Trench inner slope sediments, geochemical profiles, 1249
 Trench sediments, consolidation of, 554
 Trench slope, 494
 Tridymite lepispheres, 1274
 Turbidites, 35, 1220, 1221
 Turbidity currents, 549, 1029
 Two-ship seismic refraction data, 463, 466 Table 3, 489
 Unconformity, angular SEE: angular unconformity
 Vanadium, 1225
 Veins, 553
 defined, 576
 Velocity–depth section, 465
 Ventura Basin, 1110
 Volcanic activity, island arc, 1255
 Volcanic arc, 467
 Volcanic ash, 24, 1027, 1028, 1064, 1213
 occurrence in Leg 57 cores, 1050–1056 Table 1
 Volcanic components, 545
 Volcanic glass, 545, 1068
 alteration of, 1031
 chemical composition of, 1060, 1062–1063 Table 7
 description and refractive index, 1058
 electron microprobe analyses, 1030, 1044 Table 1
 fission track age of, 1277, 1279 Table 1
 mineralogy of, 1058
 petrography of, 1049, 1057
 refractive indices of, 1057
 settling velocity, 1057, 1060, 1065
 visual description of, 1049
 Volcanic glass shards, 1028
 chemical composition of, 1046
 refractive index of, 1043, 1044 Table 1
 Volcanic materials, 37
 diagenesis of, 550
 Volcanic rocks, AFM diagram, 1078
 conglomerates composed of, 1075
 magnetic properties of, 1078
 oxide variation diagram, 1078, 1080, Figure 5
 petrography of, 1075
 potassium–argon age, 1067, 1079
 Volcaniclastic material, 1005
 Volcanism, 1030, 1049
 explosive, 545, 1027, 1031, 1034
 in the northern Japanese islands, 1021
 late Oligocene, 4
 Volcanoes, 1058
 submarine, 1263
 Wadati-Benioff Zone, 467, 469, 482
 X-ray diffraction studies, clay minerals, 979
 methods of, 1011
 sediments, 941
 X-ray fluorescence of sediments, 1215–1216 Table 1
 Yakataga Formation, 668
 Yezo Geosyncline, 473
 Yezo Group, 474, 846, 1071, 1157
 Zeolite, burial diagenesis of, 1143, 1144, 1151
 origin of, 1149
 Zinc, 1224
 Zoophycos, 911